

The CHDP Periodicity Schedule

Screening Requirement	Age of Person Being Screened														
	< 1 mo.	1-2 mos.	3-4 mos.	5-6 mos.	7-9 mos.	10-12 mos.	13-15 mos.	16-23 mos.	2 Yr.	3 Yr.	4-5 Yr.	6-8 Yr.	9-12 Yr.	13-16 Yr.	17-20 Yr.
Interval Until Next Exam	1mo.	2 mo.	2 mos.	2 mos.	3 mos.	3 mos.	3 mos.	6 mos.	1 yr.	1 yr.	2 yr.	3 yr.	4 yr.	4 yr.	None
History & Physical Exam	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Dental Assessment	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Nutritional Assessment	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Developmental/Behavioral	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Anticipatory Guidance	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Tobacco Assesment	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Pelvic Exam 1														*	*
Measurements															
Head Circumference	*	*	*	*	*	*	*	*							
Height/Length & Weight	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Blood Pressure										*	*	*	*	*	*
Sensory Screening															
Visual Acuity Test (Snellen) 2										*	*	*	*	*	*
Clinical Observation	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Audiometric 2										*	*	*	*	*	*
Non-audiometric	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Procedures/Tests															
Tuberculin Test											*			*	
TB Exposure Risk Assessment	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hematocrit or Hemoglobin					*	*	*	*	*	*	*	*	*	*	*
Urine Dipstick or Urinalysis											*	*	*	*	*
Blood Lead Test							*	*	*	*					
Blood Lead Risk Assessment				*	*	*	*	*	*	*	*				
Other Laboratory Tests															
VDRL, RPR or ART	To be done when health history and/or physical examination warrants.														
Gonorrhea	To be done when health history and/or physical examination warrants.														
Chlamydia Test	To be done when health history and/or physical examination warrants.														
Papanicolaou (Pap) Smear	To be done when health history and/or physical examination warrants.														
Sickle Cell	To be done when health history and/or physical examination warrants.														
Ova & Parasites	To be done when health history and/or physical examination warrants.														
Immunizations	Administer as necessary to make status current.														
1. Recommended for sexually active females and females over 18 years.															
2. Test at 3 years if possible; use clinical history and observations under 3 years or if uncooperative at 3 years.															